antibodies .- online.com







anti-TRAF4 antibody (AA 437-454)



O -	4 -	D	4	page
	$T \cap$	$Pr \cap C$	шет	nane
\cup	w	1 100	IUCL	Dauc

Overview	
Quantity:	100 μg
Target:	TRAF4
Binding Specificity:	AA 437-454
Reactivity:	Human, Mouse, Rat, Cow, Guinea Pig, Horse, Rabbit, Pig, Monkey, Bat, Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAF4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))
Product Details	
Immunogen:	A synthetic peptide corresponding to a sequence at the C-terminus of human TRAF4(437-
	454aa PKFISHQDIRKRNYVRDD), identical to the related rat and mouse sequences.

Immunogen:	A synthetic peptide corresponding to a sequence at the C-terminus of human TRAF4(437-454aa PKFISHQDIRKRNYVRDD), identical to the related rat and mouse sequences.
Isotype:	IgG
Specificity:	Expressed in epithelial cells of thymus, dendritic cells of lymph node, and in the basal cell layer of epithelia such as epidermis, nasopharynx, respiratory tract, salivary gland, and esophagus.
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Purification:	Immunogen affinity purified

Target Details

Target: TRAF4

Target Details

Alternative Name:	TRAF4 (TRAF4 Products)
Background:	Name/Gene ID: TRAF4
	Synonyms: TRAF4, Malignant 62, RING finger protein 83, MLN 62, MLN62, RNF83, CART1
Gene ID:	9618

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2 HPO4, 0.05 mg Thimerosal, 0.05 mg sodium azide per 100 μ g antibody.
Preservative:	Sodium azide, Thimerosal (Merthiolate)
Precaution of Use:	This product contains Sodium azide and Thimerosal (Merthiolate): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for 1 year. After reconstitution, at 4°C for 1 month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid freeze-thaw cycles.