

Datasheet for ABIN2629892
anti-XPO1 antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	50 µL
Target:	XPO1
Binding Specificity:	AA 1025-1071, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This XPO1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), Immunofluorescence (IF), Proximity Ligation Assay (PLA)

Product Details

Brand:	IHC-plus™
Isotype:	IgG
Specificity:	Region between residue 1025 and the C-terminus (residue 1071) of human Chromosome Region Maintenance 1 (Exportin 1) using the numbering given in entry NP_003391.1 (GenelD 7514).
Purification:	Immunoaffinity purified

Target Details

Target:	XPO1
---------	------

Target Details


Alternative Name:	XP01 / CRM1 (XP01 Products)
Background:	Name/Gene ID: XP01 Synonyms: XP01, CRM1, yeast, homolog, Exp1, Exportin-1, CRM1
Gene ID:	7514
Pathways:	M Phase

Application Details

Application Notes:	Approved: IF (1:1000 - 1:10000), IHC, IHC-P (5 µg/mL), IP (2 - 10 µg/mg lysate), PLA (1:1000 - 1:10000), WB (1:2000 - 1:10000) Usage: Immunohistochemistry: Antigen retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. Human controls: Breast Carcinoma, Colon Carcinoma, Metastatic Lymph Node, Ovarian Carcinoma, Prostate Carcinoma, Testicular Seminoma.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Tris-citrate/phosphate buffer, pH 7-8, 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C for up to 1 year.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Anti-XP01 / CRM1 antibody IHC staining of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.