antibodies -online.com





anti-SERPINB6 antibody (Internal Region)





Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μg
Target:	SERPINB6
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This SERPINB6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Dumass	CEDDINIDA
Purpose:	SERPINB6
Immunogen:	Peptide with sequence C-NWDEQFDKENTEER, from the internal region of the protein sequence
	according to NP_004559.4.
Sequence:	NWDEQFDKEN TEER
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity
	chromatography using the immunizing peptide.
Grade:	Verified

Target Details

rarget betails			
Target:	SERPINB6		
Alternative Name:	SERPINB6 (SERPINB6 Products)		
Background:	SERPINB6, serpin peptidase inhibitor, clade B (ovalbumin), member 6, CAP, DKFZp686I04222, MGC111370, MSTP057, PI6, PTI, protease inhibitor 6 (placental thrombin inhibitor), serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 6		
Gene ID:	5269		
NCBI Accession:	NP_004559		
Application Details			
Application Notes:	Western Blot: Approx 40 kDa band observed in Human Ovary lysates (calculated MW of 42.6 kDa according to NP_004559.4). Recommended concentration: 0.13 µg/mL. Peptide ELISA: antibody detection limit dilution 1:128000.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	0.5 mg/mL		
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Handling Advice:	Minimize freezing and thawing.		
Storage:	-20 °C		
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.		




Image 1. ABIN263223 (0.1 μ g/ml) staining of Human Ovary lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.