

Datasheet for ABIN263730
anti-Ferritin antibody (Heavy & Light Chain)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	Ferritin (FE)
Binding Specificity:	Heavy & Light Chain
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Ferritin antibody is un-conjugated
Application:	Enzyme Immunoassay (EIA)

Product Details

Immunogen:	Source: Human Spleen
Clone:	090-10175
Isotype:	IgG2a
Characteristics:	Synonyms: Ferritin H subunit, Ferritin L subunit, FTL, FTH
Purification:	Protein A chromatography.
Purity:	> 90 % pure (SDS-PAGE)
Sterility:	0.2 µm filtered

Target Details

Target:	Ferritin (FE)
---------	---------------

Target Details

Alternative Name: Ferritin ([FE Products](#))

Background: Ferritin is a ubiquitous and highly conserved protein which plays a major role in iron homeostasis by sequestering and storing iron in a non-toxic and soluble form. It forms a holoenzyme of ~450 kDa, consisting of 24 subunits of two types, H (heavy, 21 kDa) and L (light, 19 kDa), and is capable of storing up to 4,500 atoms of ferric iron. Depending on the tissue type and physiological status of the cell, the ratio of H to L subunits in ferritin can vary widely. Ferritin is found in the liver, spleen, kidney and heart, with smaller amounts being found in blood. Serum ferritin levels serve as an indicator of the amount of iron stored in the body. Serum ferritin is the most sensitive test for anaemia, and is also used as a marker for restless leg syndrome, hemochromatosis and porphyria. As ferritin is an acute-phase reactant, it is often elevated during infection. Defects in ferritin proteins are associated with several neurodegenerative diseases. Synonyms: FTH, FTL, Ferritin H subunit, Ferritin L subunit

Pathways: [Transition Metal Ion Homeostasis](#)

Application Details

Application Notes: ELISA.
Other applications not tested.
Optimal dilutions are dependent on conditions and should be determined by the user.

Restrictions: For Research Use only

Handling

Concentration: 5.12mg/mL (OD280nm, E0.1% = 1.4)

Buffer: 10 mM Phosphate, pH 7.4 containing 150 mM Sodium chloride and 0,09 % sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C

Storage Comment: Store the antibody undiluted at 2-8 °C for one week or (in aliquots) at -40 °C for longer. If aliquoted for long term storage, fill volume should be equal to or greater than 50% of thenominal fill volume of the vial used. Avoid repeated freezing and thawing.
Shelf life: one year from despatch.

Expiry Date: 12 months