antibodies -online.com





Datasheet for ABIN2648562

G-CSF ELISA Kit



Overview

Quantity:	96 tests
Target:	G-CSF (CSF3)
Reactivity:	Human
Application:	ELISA

Product Details

Product Details	
Purpose:	Human Granulocyte Colony Stimulating Factor (G-CSF) ELISA Assay Kit (enzyme-linked immunoassay kit) is intended for the quantitative determination of human Granulocyte Colony Stimulating Factor (G-CSF) concentrations in cell culture supernates, serum, and plasma.
Sample Type:	Serum, Plasma, Cell Culture Cells
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Sensitivity:	4 pg/ml
Characteristics:	Human Granulocyte Colony Stimulating Factor (G-CSF) ELISA Assay Kit is for research use only and not to be used in diagnostic procedures.

Target Details

Target:	G-CSF (CSF3)
Alternative Name:	G-CSF (CSF3 Products)
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Actin Filament Polymerization

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Sample Volume:	100 μL
Assay Time:	3.5 h
Plate:	Pre-coated
Protocol:	Human Granulocyte Colony Stimulating Factor (G-CSF) ELISA Assay Kit employs the quantitative sandwich enzyme immunoassay technique. A monoclonal antibody specific for G-CSF has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any G-CSF present is bound by the immobilized antibody. Following incubation unbound samples are removed during a wash step, and then a detection antibody specific for G-CSF is added to the wells and binds to the combination of capture antibody- G-CSF in sample. Following a wash to remove any unbound combination, and enzyme conjugate is added to the wells. Following incubation and wash steps a substrate is added. A colored product is formed in proportion to the amount of G-CSF present in the sample. The reaction is terminated by addition of acid and absorbance is measured at 450nm. A standard curve is prepared from seven G-CSF standard dilutions and G-CSF sample concentration determined.
Restrictions:	For Research Use only
Handling	
Storage:	4 °C