

Datasheet for ABIN265483
anti-NFKBIA antibody



[Go to Product page](#)

2 Images

Overview

Quantity:	0.1 mg
Target:	NFKBIA
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFKBIA antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Specificity:	This antibody detects endogenous levels of IkappaB-alpha protein. (region surrounding Leu26)
Cross-Reactivity (Details):	Species reactivity (expected):Mouse and Rat. Species reactivity (tested):Human.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Purity:	> 95 % pure by SDS-PAGE

Target Details

Target:	NFKBIA
Alternative Name:	NFKBIA / IKBA (NFKBIA Products)
Background:	The transcription factor NFkappaB is retained in the cytoplasm in an inactive form by the

Target Details

inhibitory protein IkappaB. Activation of NFkappaB requires that IkappaB be phosphorylated on specific serine residues, which results in targeted degradation of IkappaB. IkappaB kinase alpha (IKKalpha), previously designated CHUK, interacts with IkappaB-alpha and specifically phosphorylates IkappaB-alpha on the sites that trigger its degradation Serines 32 and 36. IKKalpha appears to be critical for NFkappaB activation in response to proinflammatory cytokines. Phosphorylation of IkappaB by IKKalpha is stimulated by the NFkappaB inducing kinase (NIK), which itself is a central regulator for NFkappaB activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKKalpha, IKKbeta and IKKgamma (also designated NEMO), and each appear to make essential contributions to IkappaB phosphorylation. Synonyms: I kappa B-alpha, I-kappa-B-alpha, Ikb-alpha, IkappaBalpna, MAD3, Major histocompatibility complex enhancer-binding protein MAD3, NF-kappa-B inhibitor alpha, NFKBI

Molecular Weight:	approx. 36 kDa
Gene ID:	4792
NCBI Accession:	NP_065390
UniProt:	P25963
Pathways:	NF-kappaB Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Maintenance of Protein Location , Hepatitis C , Protein targeting to Nucleus , Toll-Like Receptors Cascades , BCR Signaling

Application Details

Application Notes:	ELISA: 1: 10000approx. 1: 40000. WB: 1: 500approx. 1: 1000. IHC: 1: 50approx. 1: 200. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only

Handling

Concentration:	1,0 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.2., 0.05 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

	should be handled by trained staff only.
Handling Advice:	DO NOT FREEZE!
Storage:	4 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C.

Images




Image 1.




Image 2.