



Datasheet for ABIN2657149  
**anti-CCR4 antibody (Alexa Fluor 647)**



[Go to Product page](#)

2 Images

### Overview

Quantity:	100 tests
Target:	CCR4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCR4 antibody is conjugated to Alexa Fluor 647
Application:	Flow Cytometry (FACS)

### Product Details

Clone:	L291H4
Isotype:	IgG1 kappa
Cross-Reactivity:	Cynomolgus
Purification:	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.

### Target Details

Target:	CCR4
Alternative Name:	CD194 ( <a href="#">CCR4 Products</a> )
Background:	CD194, also known as CCR4, is a CC chemokine receptor. It binds CCL17 and CCL22 and is expressed on a subset of T and B cells, basophils, monocytes, and NK cells. Human Th2 cells are characterized by the expression of CCR4 and CCR8, and these receptors are regulated

## Target Details

differently during Th2 development. Human peripheral blood Tregs can be divided into two distinct populations based on the expression of CCR4. Freshly isolated Tregs express CCR4 and presumably represent memory-type Tregs, and CCR4- Tregs require CD3-mediated activation to acquire a regulatory activity. Depletion of CCR4+ T cells leads to Th1-type polarization of CD4+ T cells and augmentation of CD8+ T cell responses to tumor antigens. CCR4 and its ligands are important for the recruitment of memory T cells into the skin in various cutaneous immune diseases.

## Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Buffer: Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Protect from prolonged exposure to light. Do not freeze.

Storage: 4 °C


Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C

## Images



### Flow Cytometry

#### Image 1.



## Flow Cytometry

Image 2.