

## Datasheet for ABIN2657181

# anti-CD57 antibody (Alexa Fluor 647)

# 2 Images



Go to Product page

_						
	1//	Д	rv	16	٦/	٨
U	W	$\vdash$	ΙV	Ιt	٦,	/V

Quantity:	100 tests
Target:	CD57 (B3GAT1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD57 antibody is conjugated to Alexa Fluor 647
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Blocking Reagent (BR), Co-Stimulation (CoSt)

### **Product Details**

Clone:	HNK-1
Isotype:	IgM kappa
Purification:	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.

### **Target Details**

Target:	CD57 (B3GAT1)
Alternative Name:	CD57 (B3GAT1 Products)
Background:	CD57, also known as HNK-1, NK-1, and Leu-7 is a 100-115 kD oligosaccharide antigenic determinant expressed on a variety of proteins, lipids, and chondroitin sulfate proteoglycans.  CD57 is expressed on a subset of peripheral blood lymphocytes, including NK cells and CD8+ T

#### **Target Details**

cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red blood cells, granulocytes, monocytes, or platelets. While the function of CD57 is unknown, binding to L-selectin, P-selectin, and a fragment of laminin suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).

Pathways:

Glycosaminoglycan Metabolic Process

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.		
Restrictions:	For Research Use only		
Handling			
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		


Handling Advice: Protect from prolonged exposure to light. Do not freeze.

Storage: 4 °C

Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C


should be handled by trained staff only.

#### **Images**



**Flow Cytometry** 

Image 1.



### Flow Cytometry

Image 2.