
antibodies -online.com







anti-IL21 Receptor antibody (APC)

2 Images



Go to Product page

Overview

Quantity:	100 tests
Target:	IL21 Receptor (IL21R)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This IL21 Receptor antibody is conjugated to APC
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP)

Product Details

Clone:	17A12
Isotype:	IgG1 kappa
Purification:	The antibody was purified by affinity chromatography and conjugated with APC under optimal
	conditions. The solution is free of unconjugated APC and unconjugated antibody.

Target Details

Target:	IL21 Receptor (IL21R)
Alternative Name:	CD360 (IL21R Products)
Background:	Human interleukin 21 receptor (IL-21R) is a single pass type I membrane protein and a member of the type I cytokine receptor family. Of the type I cytokine receptors, IL-21R exhibits greatest extracellular homology to the IL-2R beta subunit (i.e., it contains one copy of the WSXWS-containing cytokine-binding domain). Intracellular domains of IL-21R include the Box 1 and Box

2 elements, which are similar to the IL-9R intracellular region. Upon binding IL-21, IL-21R forms a heterodimer with the common gamma subunit (CD132) and induces Jak/Stat signaling. IL-21R is expressed on B cells and at various levels on NK and T cells. IL-21 is a potent immunomodulatory cytokine mainly produced by NKT and CD4 T-cells (particularly the inflammatory Th17 subset) and has pleiotropic effects on both innate and adaptive immune responses. These actions include positive effects such as enhanced proliferation of natural killer (NK) cells and cytotoxic T cells that can destroy virally infected or cancerous cells and direct inhibitory effects on the antigen-presenting function of dendritic cells, and can be proapoptotic for B cells and NK cells. Studies have shown that IL-21 is also an autocrine cytokine that potently induces Th17 differentiation and suppresses Foxp3 expression, and serves as a target for treating inflammatory diseases.

Pathways:


JAK-STAT Signaling

Application Details

Application Notes:


Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4°C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Optimal working dilution should be determined by the investigator.



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.