

Datasheet for ABIN2658299 anti-CX3CR1 antibody (APC)

2 Images



Overview

| Quantity: | 100 tests |
|--------------|---|
| Target: | CX3CR1 |
| Reactivity: | Human |
| Host: | Rat |
| Clonality: | Monoclonal |
| Conjugate: | This CX3CR1 antibody is conjugated to APC |
| Application: | Biochemical Assay (BCA) |

Product Details

| Clone: | 2A9-1 |
|---------------|--|
| Isotype: | IgG2b kappa |
| Purification: | The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody. |

Target Details

| Target: | CX3CR1 |
|-------------------|---|
| Alternative Name: | CX3CR1 (CX3CR1 Products) |
| Background: | CX3CR1 is a G-protein-coupled seven-transmembrane chemokine receptor, also called GPR13 |
| | or V28. It is expressed on NK cells, T cell subset, monocytes/macrophages, dendritic cells, and |
| | some malignant epithelial cells. CX3CL1 (known also as fractalkine and neurotactin) is the |
| | ligand of CX3CR1. CX3CL1 is a unique transmembrane molecule with a CX3C-motif chemokine |

Target Details

| domain and a mucin-like stalk. CX3CL1 is expressed by activated-endothelial cells, neurons | , and |
|--|-------|
| astrocytes. The interaction of CX3CR1 and its ligand mediates firm cell adhesion and migra | tion. |

Pathways:

Cellular Response to Molecule of Bacterial Origin


Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling


| Buffer: | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA . |
|--------------------|--|
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Protect from prolonged exposure to light. Do not freeze. |
| Storage: | 4 °C |
| Storage Comment: | The antibody solution should be stored undiluted between 2°C and 8°C. |

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.