

Datasheet for ABIN2658649  
**anti-CD39 antibody (APC)**[Go to Product page](#)

## 2 Images

## Overview

|              |   |
|--------------|---|
| Quantity:    | 100 tests                               |
| Target:      | CD39 (ENTPD1)                           |
| Reactivity:  | Human                                   |
| Host:        | Mouse                                   |
| Clonality:   | Monoclonal                              |
| Conjugate:   | This CD39 antibody is conjugated to APC |
| Application: | Biochemical Assay (BCA)                 |

## Product Details

|                   |   |
|-------------------|---|
| Clone:            | A1  |
| Isotype:          | IgG1 kappa  |
| Cross-Reactivity: | Rhesus Monkey   |
| Purification:     | The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody. |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | CD39 (ENTPD1)  |
| Alternative Name: | CD39 ( <a href="#">ENTPD1 Products</a> )   |
| Background:       | Human CD39 is an integral membrane protein with two transmembrane domains. It exists as a homotetramer. Expression of CD39 is found on activated lymphocytes, a subset of T cells and B cells, and dendritic cells with weak staining on monocytes and granulocytes. CD39 and CD73 |

Target Details

have been found on regulatory T cells, specifically the effector/memory like T cells. CD39 can hydrolyze both nucleoside triphosphates and diphosphates. CD39 is the dominant ecto nucleotidase of vascular and placental trophoblastic tissues and appears to modulate the functional expression of type 2 purinergic (P2) G protein coupled receptors (GPCRs). CD39 has intrinsic ecto-ATPase activity. Expression of CD39 is induced on T cells and increased on B cells as a late activation antigen.


Application Details

|                    |  |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions:      | For Research Use only  |

Handling


|                    |  |
|--------------------|--|
| Buffer:            | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .                              |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice:   | Protect from prolonged exposure to light. Do not freeze.   |
| Storage:           | 4 °C   |
| Storage Comment:   | The antibody solution should be stored undiluted between 2°C and 8°C.  |

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.