

Datasheet for ABIN2658679
anti-ITGA4 antibody (APC)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 tests
Target:	ITGA4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ITGA4 antibody is conjugated to APC
Application:	Flow Cytometry (FACS), Intracellular Flow Cytometry (ICFC)

Product Details

Clone:	9F10
Isotype:	IgG1 kappa
Cross-Reactivity:	Baboon, Chimpanzee, Marmoset, Cynomolgus, Rhesus Monkey, Squirrel Monkey (Saimiri spec.), Horse (Equine), Cow (Bovine), Sheep (Ovine), Dog (Canine), Cat (Feline)
Purification:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Target Details

Target:	ITGA4
Alternative Name:	CD49d (ITGA4 Products)
Background:	CD49d is a 150 kD α integrin chain known as $\alpha 4$ integrin or VLA-4 α chain. It forms a heterodimer with either integrin $\beta 1$ ($\alpha 4 \beta 1$, VLA-4) or $\beta 7$ ($\alpha 4 \beta 7$). CD49d is expressed broadly on

Target Details

T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin. $\alpha 4\beta 7$ is the receptor for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.

Pathways: [Integrin Complex](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Buffer: Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Protect from prolonged exposure to light. Do not freeze.

Storage: 4 °C

Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C.



Flow Cytometry

Image 1.