

Datasheet for ABIN2658695

anti-CD56 antibody (APC)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
U	W	\vdash	ΙV	Ιt	٦,	/V

Quantity:	100 tests	
Target:	CD56 (NCAM1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD56 antibody is conjugated to APC	
Application:	Flow Cytometry (FACS), Intracellular Flow Cytometry (ICFC)	

Product Details

Clone:	MEM-188
Isotype:	IgG2a kappa
Cross-Reactivity:	Cow (Bovine), Pig (Porcine)
Purification:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.


Target Details

Target:	CD56 (NCAM1)
Alternative Name:	CD56 (NCAM1 Products)
Background:	CD56 is a single transmembrane glycoprotein also known as N-CAM (Neural Cell Adhesion Molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is
	expressed on NK cells and NK-T cells. CD56 is also expressed in brain (cerebellum and cortex)

and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparin sulfate.

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.
Images	



Flow Cytometry

Image 1.