

## Datasheet for ABIN2658700 anti-L-Selectin antibody (APC)

### 1 Image



Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	100 tests
Target:	L-Selectin (SELL)
Reactivity:	Chemical
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This L-Selectin antibody is conjugated to APC
Application:	Flow Cytometry (FACS), Intracellular Flow Cytometry (ICFC)

#### **Product Details**

Clone:	DREG-56	
Isotype:	IgG1 kappa	
Cross-Reactivity:	Chimpanzee, Cow (Bovine)	
Purification:	The antibody was purified by affinity chromatography, and conjugated with APC under optima conditions. The solution is free of unconjugated APC and unconjugated antibody.	

#### Target Details


Target:	L-Selectin (SELL)	
Alternative Name:	CD62L (SELL Products)	
Target Type:	Chemical	
Background:	CD62L is a 74-95 kD single chain type I glycoprotein referred to as L-selectin or LECAM-1. It is	

expressed on most peripheral blood B cells, subsets of T and NK cells, monocytes, granulocytes, and certain hematopoietic malignant cells. CD62L binds to carbohydrates present on certain glycoforms of CD34, glycam-1, and MAdCAM-1 and with a low affinity to anionic oligosaccharide sequences related to sialylated Lewis X (sLex, CD15s) through its C-type lectin domain. CD62L is important for the homing of naive lymphocytes to high endothelial venules in peripheral lymph nodes and Peyer's patches. It also plays a role in leukocyte rolling on activated endothelial cells.

#### **Application Details**

**Images** 

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.



# Flow Cytometry Image 1.