

Datasheet for ABIN2658799

anti-Integrin beta 7 antibody (APC)





\sim			
	ve	r\/	٨
\cup	V C	1 V I	٧V

Overview	
Quantity:	100 tests
Target:	Integrin beta 7 (ITGB7)
Reactivity:	Human, Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Integrin beta 7 antibody is conjugated to APC
Application:	Flow Cytometry (FACS), Intracellular Flow Cytometry (ICFC)
Product Details	
Clone:	FIB504

Clone:	FIB504	
Isotype:	IgG2a kappa	
Cross-Reactivity:	Chimpanzee, Baboon, Cynomolgus, Rhesus Monkey	
Purification:	The antibody was purified by affinity chromatography, and conjugated with APC under optimic conditions. The solution is free of unconjugated APC and unconjugated antibody.	

Target Details

Target:	Integrin beta 7 (ITGB7)	
Alternative Name:	Integrin Beta7 (ITGB7 Products)	
Background:	Integrin β 7 is a 130 kD glycoprotein also known as integin β p. It is a member of the Ig superfamily. In association with integrin α 4 or α E chain, β 7 forms α 4/ β 7 or α E/ β 7 heterodimer. α 4/ β 7 (CD49d/ β 7, LPAM-1) is expressed on the majority of peripheral lymphocytes, on small	

Target Details

subsets of thymocytes, and bone marrow progenitors. LPAM-1 binds to several ligands, VCAM-1, MAdCAM-1 and fibronectin, and is involved in lymphocyte adhesion and some hematopoietic progenitor cells migration. $\alpha E/\beta T$ (CD103/ βT , $\alpha IEL/\beta T$) is expressed on intestinal intraepithelial lymphocytes (IEL), dendritic epidermal T cells, T regulatory cells, a subset of CD8+ T cells in lymph nodes and lamina propria. CD103/ βT complex is thought to play a role in lymphocyte retention via interaction with its ligand E-Cadherin.


Pathways:

Integrin Complex

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.