

Datasheet for ABIN2660016
anti-CCR4 antibody (PerCP-Cy5.5)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 tests
Target:	CCR4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCR4 antibody is conjugated to PerCP-Cy5.5
Application:	ELISA (Detection)

Product Details

Clone:	L291H4
Isotype:	IgG1 kappa
Cross-Reactivity:	Cynomolgus
Purification:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Target Details

Target:	CCR4
Alternative Name:	CD194 (CCR4 Products)
Background:	CD194, also known as CCR4, is a CC chemokine receptor. It binds CCL17 and CCL22 and is expressed on a subset of T and B cells, basophils, monocytes, and NK cells. Human Th2 cells

Target Details


are characterized by the expression of CCR4 and CCR8, and these receptors are regulated differently during Th2 development. Human peripheral blood Tregs can be divided into two distinct populations based on the expression of CCR4. Freshly isolated Tregs express CCR4 and presumably represent memory-type Tregs, and CCR4- Tregs require CD3-mediated activation to acquire a regulatory activity. Depletion of CCR4+ T cells leads to Th1-type polarization of CD4+ T cells and augmentation of CD8+ T cell responses to tumor antigens. CCR4 and its ligands are important for the recruitment of memory T cells into the skin in various cutaneous immune diseases.

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only


Handling

Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.