
antibodies - online.com







anti-CCR6 antibody (PerCP-Cy5.5)





Overview

Quantity:	100 tests
Target:	CCR6
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCR6 antibody is conjugated to PerCP-Cy5.5
Application:	Immunofluorescence (IF)

Product Details

Clone:	G034E3
Isotype:	IgG2b kappa
Cross-Reactivity:	Cynomolgus, Rhesus Monkey
Purification:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Target Details

Target:	CCR6
Alternative Name:	CD196 (CCR6 Products)
Background:	CCR6, also known as CD196, is a chemokine receptor that is expressed on immature dendritic
	cells, B lymphocytes, and memory T cells. CCR6 binds CCL20, although members of the $\boldsymbol{\beta}$

Target Details

defensin family also bind CCR6 with a lower affinity. CCR6 positive cells, and its ligand CCL20, have been detected in numerous organs, especially the secondary lymphoid organ. CCL20 is selectively made by the follicle-associated epithelium (FAE) overlying Peyers patches (PPs) and isolated lymphoid follicles (ILFs). CCL20 contributes to the recruitment of CCR6-expressing B cells to these structures. In humans, CCR6 can function to mediate arrest of T cells on dermal endothelial cells and is highly expressed on T cells resident in both normal and psoriatic skin. CCR6 and/or CCL20 have been implicated in the pathogenesis of rheumatoid arthritis and inflammatory bowel disease. Human T cells that are able to produce IL-17 express CCR6. It suggests that CCL20 and CCR6 have a role in inflammatory diseases by recruiting Th17 cells to target tissues.

Pathways:


cAMP Metabolic Process

Application Details

Application Notes:


Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4°C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Optimal working dilution should be determined by the investigator.



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.