

Datasheet for ABIN2660079

anti-Neuropilin 1 antibody (PerCP-Cy5.5)[Go to Product page](#)**2** Images

Overview

Quantity:	100 µg
Target:	Neuropilin 1 (NRP1)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Neuropilin 1 antibody is conjugated to PerCP-Cy5.5
Application:	Flow Cytometry (FACS)

Product Details

Clone:	3E12
Isotype:	IgG2a kappa
Purification:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Target Details

Target:	Neuropilin 1 (NRP1)
Alternative Name:	CD304 (NRP1 Products)
Background:	CD304, also known as neuropilin-1, is a 140 kD type I transmembrane protein. Its extracellular region contains two CUB, two FV/FVIII, and one MAM domain. It is expressed by natural regulatory T cells (nTreg), a subset of invariant natural killer T cells (iNKT), endothelial cells, and

Target Details

neurons. Neuropilin-1 stabilizes the interaction between Tregs and dendritic cells, contributes to the recruitment of tumor-infiltrating Tregs in response to tumor-derived VEGF, and influences the process of angiogenesis and axon guidance. The ligands of CD304 are VEGF165 and semaphorin-3A.

Pathways: [Regulation of Cell Size](#), [Signaling Events mediated by VEGFR1 and VEGFR2](#), [Smooth Muscle Cell Migration](#), [Platelet-derived growth Factor Receptor Signaling](#), [VEGFR1 Specific Signals](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Concentration: 0.2 mg/mL

Buffer: Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Protect from prolonged exposure to light. Do not freeze.

Storage: 4 °C


Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.