

Datasheet for ABIN2660215

anti-Integrin beta 7 antibody (PerCP-Cy5.5)





Overview

Quantity:	100 tests
Target:	Integrin beta 7 (ITGB7)
Reactivity:	Human, Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Integrin beta 7 antibody is conjugated to PerCP-Cy5.5
Application:	Flow Cytometry (FACS)

Product Details

Clone:	FIB504
Isotype:	IgG2a kappa
Cross-Reactivity:	Chimpanzee, Baboon, Cynomolgus, Rhesus Monkey
Purification:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Target Details

Target:	Integrin beta 7 (ITGB7)
Alternative Name:	Integrin Beta7 (ITGB7 Products)
Background:	Integrin $\beta 7$ is a 130 kD glycoprotein also known as integin βp . It is a member of the Ig
	superfamily. In association with integrin $\alpha 4$ or αE chain, $\beta 7$ forms $\alpha 4/\beta 7$ or $\alpha E/\beta 7$ heterodimer.

Target Details

 $\alpha4/\beta7$ (CD49d/ $\beta7$, LPAM-1) is expressed on the majority of peripheral lymphocytes, on small subsets of thymocytes, and bone marrow progenitors. LPAM-1 binds to several ligands, VCAM-1, MAdCAM-1 and fibronectin, and is involved in lymphocyte adhesion and some hematopoietic progenitor cells migration. $\alpha E/\beta7$ (CD103/ $\beta7$, α IEL/ $\beta7$) is expressed on intestinal intraepithelial lymphocytes (IEL), dendritic epidermal T cells, T regulatory cells, a subset of CD8+ T cells in lymph nodes and lamina propria. CD103/ $\beta7$ complex is thought to play a role in lymphocyte retention via interaction with its ligand E-Cadherin.

Pathways:

Integrin Complex

4°C

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Buffer: Preservative:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA . Sodium azide


Protect from prolonged exposure to light. Do not freeze.

Storage Comment:

Handling Advice:

Storage:

The antibody solution should be stored undiluted between 2°C and 8°C.



Flow Cytometry

Image 1.