

Datasheet for ABIN2660598

anti-CD56 antibody (APC-Cy7)

2 Images



Go to Product page

Overview

Quantity:	100 tests
Target:	CD56 (NCAM1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD56 antibody is conjugated to APC-Cy7
Application:	Biochemical Assay (BCA)

Product Details

Clone:	HCD56
Isotype:	IgG1 kappa
Purification:	The antibody was purified by affinity chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

Target Details


l arget:	CD56 (NCAMT)
Alternative Name:	CD56 (NCAM1 Products)
Background:	CD56 is a single transmembrane glycoprotein also known as NCAM (Neural Cell Adhesion
	Molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is
	expressed on NK cells and NK-T cells. CD56 is also expressed in the brain (cerebellum and
	cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias,

small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparin sulfate.

Application Details


Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
riarraming	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.