# antibodies -online.com







Image



Go to Product page

_						
( )	V	ρ	rv	16	$\supset I$	Λ

Quantity:	50 μg
Target:	СЗа
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This C3a antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS)

### **Product Details**

Clone:	D17-1
Isotype:	IgG1 kappa
Purification:	The antibody was purified by affinity chromatography, and conjugated with biotin under optimal
	conditions. The solution is free of unconjugated biotin.

# Target Details

Target:	C3a
Alternative Name:	C3a/C3a/C3 (C3a Products)
Background:	C3, complement component 3, is a 195 KD glycoprotein heterodimer that is linked by disulfide bonds. It is the fourth complement component to react in the classical pathway and it is also a
	reactant in the alternative complement pathway. C3 convertase splits C3 into C3b and C3a. C3b
	can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune

#### **Target Details**

aggregates. Macrophages and neutrophils have receptors for C3b. C3a is released into the surrounding fluids. It can bind to receptors on basophils and mast cells triggering them to release their vasoactive contents (e.g., histamine). Because of the role of these materials in anaphylaxis, C3a is called an anaphylatoxin. In blood, the C3a fragment is rapidly desarginated by serum carboxypeptidase N enzyme to generate C3a desArg. Both C3a and C3a desArg are not glycosylated.

Pathways:


Complement System

# **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

# Handling

Concentration:	0.5 mg/mL	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.	
Storage:	4 °C	
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.	



#### **ELISA**

Image 1.