

Datasheet for ABIN2660920

anti-CD11c antibody (Biotin)

1 Image



Overview

| Quantity: | 100 μg |
|--------------|---|
| Target: | CD11c (ITGAX) |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This CD11c antibody is conjugated to Biotin |
| Application: | Biochemical Assay (BCA) |

Product Details

| Clone: | 3-9 |
|-------------------|---|
| Isotype: | IgG1 kappa |
| Cross-Reactivity: | Cynomolgus, Rhesus Monkey, Chlorocebus aethiops (Green monkey) (Cercopithecus aethiops), Baboon, Chimpanzee, Squirrel Monkey (Saimiri spec.), Green Monkey (Chlorocebus sabaeus) |
| Purification: | The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin. |

Target Details

| Target: | CD11c (ITGAX) |
|-------------------|--|
| Alternative Name: | CD11c (ITGAX Products) |
| Background: | CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin αX and CR4. |
| | CD11c non-covalently associates with integrin $\beta 2$ (CD18) and is expressed on |

Target Details

| monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T $\mbox{\ensuremath{a}}$ | nd B cells. |
|--|-------------|
| CD11c has been reported to play a role in adhesion and CTL killing through its intera | ctions with |
| fibrinogen, CD54, and iC3b. | |

Pathways:

Complement System, Activated T Cell Proliferation, Integrin Complex


Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Concentration: | 0.5 mg/mL |
|--------------------|--|
| Buffer: | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Do not freeze. |
| Storage: | 4 °C |
| Storage Comment: | The antibody solution should be stored undiluted between 2°C and 8°C. |

Images



Flow Cytometry

Image 1.