

Datasheet for ABIN2661108

anti-CD51 antibody (Biotin)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
U	W	\vdash	ΙV	Ιt	٦,	/V

Quantity:	50 μg
Target:	CD51 (ITGAV)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CD51 antibody is conjugated to Biotin
Application:	Immunofluorescence (IF), Intracellular Flow Cytometry (ICFC)

Product Details

Clone:	RMV-7
Isotype:	IgG1 kappa
Purification:	The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Target Details

Target:	CD51 (ITGAV)
Alternative Name:	CD51 (ITGAV Products)
Background:	CD51 is a 140 kD protein, also known as αV integrin, vitronectin receptor, and integrin αV . It is a member of the integrin family, expressed on activated T cells, polymorphonuclear granulocytes, platelets, blastocysts, and osteoclasts. CD51 forms heterodimers by association with integrins $\beta 1$, $\beta 3$, $\beta 5$ or $\beta 6$, these complexes then act as receptors for multiple extracellular matrix

Target Details

proteins (ECM). The av integrin heterodimers have varied functions in development, stimulation/activation and homeostasis. The primary ligands for CD51 complexes are fibronectin, fibrinogen, vitronectin, thrombspondin, von Willebrand factor, and CD31. The RMV-7 antibody has been reported to block binding of CD51 to vitronectin, fibronectin, and CD31 in some cell types, as well as blocking LAK cell cytotoxicity.

Pathways:


Cell-Cell Junction Organization, Signaling Events mediated by VEGFR1 and VEGFR2, Growth Factor Binding, Integrin Complex

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.



Flow Cytometry

Image 1.