

Datasheet for ABIN2661741 anti-CD45RB antibody (FITC)

Image



| _ | | | | |
|---|----|-----|-----|---|
| | ve | rVI | 161 | M |

| Quantity: | 50 μg |
|--------------|--|
| Target: | CD45RB |
| Reactivity: | Mouse |
| Host: | Rat |
| Clonality: | Monoclonal |
| Conjugate: | This CD45RB antibody is conjugated to FITC |
| Application: | Flow Cytometry (FACS), Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP) |

Product Details

| Clone: | C363-16A |
|---------------|---|
| Isotype: | IgG2a kappa |
| Purification: | The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC. |

Target Details

| Target: | CD45RB |
|-------------------|--|
| Alternative Name: | CD45RB (CD45RB Products) |
| Background: | CD45RB is an isoform of CD45 with exon 5 splicing (encodes B determinant). It is a 220 kD glycoprotein expressed on peripheral B cells, naive T cells, thymocytes, weakly on |
| | macrophages, and dendritic cells. It plays a critical role in TCR and BCR signaling. As T cells |

become activated and progress from naive to memory cells, CD45RB expression is downregulated. Additionally, functionally distinct CD4+ T cell subsets, which secrete differing cytokine profiles, can be separated by CD45RB intensity. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4 and Thy-1.


Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Concentration: | 0.5 mg/mL | |
|--------------------|--|--|
| Buffer: | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Handling Advice: | Protect from prolonged exposure to light. Do not freeze. | |
| Storage: | 4 °C | |
| Storage Comment: | The antibody solution should be stored undiluted between 2°C and 8°C. | |

Images



Flow Cytometry

Image 1.