

Datasheet for ABIN2661878
anti-ESAM antibody (FITC)



[Go to Product page](#)

1 Image

Overview

Quantity:	50 µg
Target:	ESAM
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This ESAM antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Flow Cytometry (FACS), Blocking Reagent (BR)

Product Details

Clone:	1G8-ESAM
Isotype:	IgG2a kappa
Purification:	The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Target Details

Target:	ESAM
Alternative Name:	ESAM (ESAM Products)
Background:	Endothelial cell-selective adhesion molecule (ESAM) is a 55-kD membrane protein composed of two extracellular Ig domains, a single transmembrane domain, and a cytoplasmic domain. ESAM is predominantly expressed at endothelial junctions and on platelets, participating in the

Target Details

migration of neutrophils through the vessel wall by influencing endothelial cell contacts. It impacts vascular permeability and extravasation. Recently, it was reported that ESAM is a novel marker for murine hematopoietic stem cells (HSCs) in fetal liver. ESAM expression is correlated with HSC activity. The ESAM^{Hi} population was highly enriched for multipotent myeloid-erythroid progenitors and primitive progenitors with lymphopoietic activity, and exclusively reconstituted long-term lymphohematopoiesis in lethally irradiated recipients.


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.