# antibodies -online.com









### Overview

Quantity:	100 tests
Target:	IKZF2
Reactivity:	Human, Mouse
Host:	Armenian Hamster
Clonality:	Monoclonal
Conjugate:	This IKZF2 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

# Product Details

Clone:	22F6
Isotype:	IgG
Purification:	The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

# Target Details


Target:	IKZF2
Alternative Name:	Helios (IKZF2 Products)
Background:	Helios is a member of the Ikaros family of zinc finger transcription factors. It contains a C-terminal region composed of 2 zinc-finger domains that mediate dimerization between the
	family members. Helios was originally cloned from a mouse thymoma line. It is mainly
	expressed in peripheral T cells and thymocytes. It is found at high levels in a subpopulation of

regulatory T cells. Helios plays an important role in T cell development and homeostasis. Overexpression of Helios profoundly alters  $\alpha\beta$  T cell differentiation and activation. It has been determined that silencing of Helios in B cells is critical for maintaining normal B cell function. Helios is also involved in tumor immunity.

### **Application Details**


Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide and 0.2 % (w/v) BSA .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

### **Images**



# **Flow Cytometry**

Image 1.



## Flow Cytometry

Image 2.