

# Datasheet for ABIN2662954 anti-CSF1R antibody (PE)

## 2 Images



Go to Product page

#### Overview

| Quantity:         | 50 μg   |
|-------------------|---|
| Target:           | CSF1R   |
| Reactivity:       | Mouse   |
| Host:             | Rat   |
| Clonality:        | Monoclonal  |
| Conjugate:        | This CSF1R antibody is conjugated to PE   |
| Application:      | Immunohistochemistry (IHC), Flow Cytometry (FACS), Immunofluorescence (IF)  |
| Product Details   |   |
| Clone:            | AFS98   |
| Isotype:          | IgG2a kappa   |
| Purification:     | The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.   |
| Target Details    |   |
| Target:           | CSF1R   |
| Alternative Name: | CD115 (CSF1R Products)  |
| Background:       | CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have |

described an Ig-like extracellular domain, a transmembrane domain, an intracellular

#### **Target Details**

juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

Pathways:

RTK Signaling, Inositol Metabolic Process, Cell-Cell Junction Organization


#### **Application Details**

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions:      | For Research Use only  |

#### Handling


| Concentration:     | 0.2 mg/mL  |
|--------------------|--|
| Buffer:            | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.   |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice:   | Protect from prolonged exposure to light. Do not freeze.   |
| Storage:           | 4 °C   |
| Storage Comment:   | The antibody solution should be stored undiluted between 2°C and 8°C.  |

#### **Images**



#### **Flow Cytometry**

Image 1.



### Flow Cytometry

Image 2.