Datasheet for ABIN2663504

anti-ITGA4 antibody (PE)

1 Image

Go to Product page

Overview

Quantity:	50 µg
Target:	ITGA4
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This ITGA4 antibody is conjugated to PE
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF)

Product Details

Clone:	9C10 (MFR4-B)
Isotype:	IgG2a kappa
Purification:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Target Details

Target:	ITGA4
Alternative Name:	CD49d (ITGA4 Products)
Background:	CD49d is a 150 kD glycoprotein, also known as α4 integrin or VLA-4 α chain. It is a member of the integrin family, expressed on T and B cells, monocytes, eosinophils, basophils, mast cells, thymocytes, NK cells, and dendritic cells. CD49d is a heterodimer expressed with either of two β

Target Details

	chains, β1 (CD29) or β7, to form the VLA-4 (integrin α4β1) or LPAM-1 (integrin α4β7) complexes. CD49d plays a critical role in both adhesion and T cell costimulation. The primary ligands for CD49d are VCAM-1, MAdCAM-1, and fibronectin. The 9C10(MFR4.B) antibody, in combination with the R1-2 monoclonal antibody, completely blocks VCAM-1 binding to VLA-4.
Pathways:	Integrin Complex

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.2 mg/mL
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from prolonged exposure to light. Do not freeze.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.