antibodies -online.com









Go to Product page

_					
U	V	er	VI	е	W

Quantity:	50 μg
Target:	CD93
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CD93 antibody is conjugated to PE
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Clone:	AA4-1
Isotype:	lgG2b kappa
Purification:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal
	conditions. The solution is free of unconjugated PE and unconjugated antibody.

Target Details

Target:	CD93
Alternative Name:	CD93 (CD93 Products)
Target Type:	Chemical
Background:	CD93 is a 130-140kD C-type lectin like type I transmembrane protein, also known as complement component 1, q subcomponent (C1qR1), C1qRp collectin receptor (C1qRp), or AA4 antigen. It is a receptor expressed on immature B lymphocytes, hematopoietic progenitors

and stem cells in adult bone marrow, fetal liver, embryonic yolk sac. CD93 expression level on splenic immature/transitional B cells is much lower than in bone marrow. It is reinduced during plasma cell differentiation and plays an important role in maintaining plasma cells in bone marrow niches. Immature dendritic cells express CD93 and down-regulate this molecule upon maturation, suggesting a role in uptake of particles by DC. It is also expressed on monocytes, macrophages, and endothelial cells. Macrophages from CD93 (-/-) mice had a significant phagocytic defect in the clearance of apoptotic cells in vivo, indicating CD93 may contribute to the in vivo clearance of dying cells. Binding of CD93 to C1q remains controversial.

Application Details

Handling Advice:


Storage:

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Concentration:	0.2 mg/mL	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	

Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C.


4°C

Protect from prolonged exposure to light. Do not freeze.



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.