

Datasheet for ABIN2663819 anti-ITGA4 antibody (PE)





Go to Product page

| \sim | | | | |
|--------|-----|-------|------------|----|
| | Ive | r\/I | $ \Delta $ | Λ/ |
| | V C | 1 V I | | vv |

| Quantity: | 50 μg |
|--------------|---|
| Target: | ITGA4 |
| Reactivity: | Mouse |
| Host: | Rat |
| Clonality: | Monoclonal |
| Conjugate: | This ITGA4 antibody is conjugated to PE |
| Application: | ELISA (Capture) |

Product Details

| Clone: | DATK32 |
|---------------|--|
| Isotype: | IgG2a kappa |
| Purification: | The antibody was purified by affinity chromatography, and conjugated with PE under optimal |
| | conditions. The solution is free of unconjugated PE and unconjugated antibody. |

Target Details

| Target: | ITGA4 | |
|-------------------|---|--|
| Alternative Name: | LPAM-1 (ITGA4 Products) | |
| Background: | DATK32 antibody is specific for a combinatorial determinate of integrin $\alpha 4 \beta 7$ complex. Integrin $\alpha 4\beta 7$ is composed of a 150 kD ($\alpha 4$ or CD49d) and a 130 kD ($\beta 7$) heterodimer, also known as | |
| | CD49d/β7 or LPAM-1. Belonging to the Ig superfamily, it is found on the majority of peripheral | |
| | lymphocytes and subsets of thymocytes and bone marrow cells (including mast cell | |

Target Details

| progenitors). Integrin $\alpha 4\beta 7$ binds its ligands, VCAM-1 (CD106), MAdCAM-1 and fibronectin, and |
|---|
| plays an important role in lymphocytes adhesion and the direction of migration of blood |
| lymphocytes to the intestine and associated lymphoid tissues. |

Pathways:

Integrin Complex


Application Details

| Application Notes: Optimal working dilution should be determined by the investigator. | |
|---|-----------------------|
| Restrictions: | For Research Use only |

Handling


| Concentration: | 0.2 mg/mL | |
|--------------------|--|--|
| Buffer: | Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Handling Advice: | Protect from prolonged exposure to light. Do not freeze. | |
| Storage: | 4 °C | |
| Storage Comment: | The antibody solution should be stored undiluted between 2°C and 8°C. | |

Images




Flow Cytometry

Image 1.



Flow Cytometry

Image 2.



Flow Cytometry

Image 3.