

Datasheet for ABIN2664097
anti-CCL28 antibody (Biotin)[Go to Product page](#)

1 Image

Overview

Quantity:	50 µg
Target:	CCL28
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCL28 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Clone:	J139G11
Isotype:	IgG2b kappa
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	CCL28
Alternative Name:	CCL28 (CCL28 Products)
Background:	Human CCL28 was identified in an EST database using a human chemokine consensus sequence. CCL28 is constitutively expressed by mucosal epithelial cells such as exocrine glands, trachea, and colon. Acinar epithelial cells in human and mouse salivary glands are strongly positive for CCL28 by immunostaining, and human saliva and milk contain high concentrations of CCL28. Furthermore, it is known that human and mouse CCL28 have

Target Details

antimicrobial activity attributed to the C-terminus of CCL28 against *Candida albicans*, Gram-negative bacteria, and Gram-positive bacteria. In addition, using a CCR10 deficient mouse model, it was shown that CCR10 plays a critical role in localization and accumulation of IgA ASC in lactating mammary glands. Also, it has been described that CCL28 is upregulated in epithelial inflammation, and this upregulation allows the recruitment of Tregs expressing CCR10. In fact, a subpopulation of CCR10-expressing CD25+CD4+FOXP3+ Tregs with potent anti-inflammatory properties was isolated from a chronically inflamed human liver. Information suggests that tumor hypoxia promotes the recruitment of Tregs through induction of expression of CCL28, and this induction is mediated by the hypoxia inducible factor-1 α (HIF-1 α). Therefore, tumor hypoxia promotes tolerance and angiogenesis via CCL28 and Tregs.

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Concentration:	0.5 mg/mL
----------------	-----------


Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.
---------	--

Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
--------------------	--

Storage:	4 °C
----------	------

Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.
------------------	---



ELISA

Image 1.