antibodies - online.com





anti-CD207 antibody

Images



_						
()	V	ρ	rv	16	\supset	Λ

Quantity:	100 μg
Target:	CD207
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD207 antibody is un-conjugated
Application:	Intracellular Flow Cytometry (ICFC)

Product Details

Clone:	10E2
Isotype:	IgG1 kappa
Cross-Reactivity:	Cynomolgus
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	CD207
Alternative Name:	CD207 (CD207 Products)
Background:	CD207, also known as Langerin, is a 40 kD type II transmemberane cell glycoprotein which
	belongs to C-type lectin with mannose binding specificity. It is predominantly expressed on
	Langerhans cells and induces the formation of Birbeck granules, the Langerhans cell hallmark
	organelle. It is also found on several other subtypes of dendritic cells, such as dermal CD103-

positive dendritic cells and splenic CD8-positive dendritic cells. Langerin is generally thought to be involved in antigen processing. Recently, it has been found that HIV captured by Langerin was internalized into Birbeck granule and degraded, which results in inhibition of HIV-1 infection and subsequent transmission.


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	

Handling


Concentration:	0.5 mg/mL	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.	

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.