

Datasheet for ABIN2664469

anti-HVEM antibody**1** Image[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HVEM (TNFRSF14)
Reactivity:	Mouse
Host:	Armenian Hamster
Clonality:	Monoclonal
Conjugate:	This HVEM antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Flow Cytometry (FACS), Blocking Reagent (BR)

Product Details

Clone:	HMHV-1B18
Isotype:	IgG
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	HVEM (TNFRSF14)
Alternative Name:	CD270 (TNFRSF14 Products)
Target Type:	Viral Protein
Background:	Herpes Virus Entry Mediator (HVEM, TR2) is a type I transmembrane protein of TNF-receptor superfamily. This receptor was identified as a cellular mediator of herpes simplex virus (HAS) entry. Binding of HSV viral envelope glycoprotein D to this receptor has been shown to be part of the viral entry mechanism. It is expressed on most cell types, including T cells, B cells,

Target Details

monocytes, neutrophils, and dendritic cells. It is also found in brain, heart, kidney, liver, and other organs. The ligands of HVEM are LIGHT, BTLA, LT α , and CD160. HVEM activates NF- κ B through the TNF-related cytokine LIGHT to serve as a costimulatory pathway during T cell activation. HVEM also functions as a ligand for the Ig superfamily members B and T lymphocyte attenuator (BTLA) and CD160 to deliver a coinhibitory signal and limit inflammatory responses initiated by T cells. HVEM plays an important role in regulating lymphocyte activation and homeostasis in immune responses.

Pathways: [Production of Molecular Mediator of Immune Response, Cancer Immune Checkpoints](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Concentration: 0.5 mg/mL


Buffer: Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C.



Flow Cytometry

Image 1.