

Datasheet for ABIN2664561

anti-BST2 antibody

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	BST2
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This BST2 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Blocking Reagent (BR)
Product Details	
Clone:	129c1
Isotype:	IgG2b kappa

Purification:

The antibody was purified by affinity chromatography.

Target Details

Target:	BST2
Alternative Name:	CD317 (BST2 Products)
Background:	CD317, known as BST2, tetherin, HM1.2 antigen, bone marrow stromal antigen 2, or PDCA-1, is type II transmembrane glycoprotein with a molecular mass of 29-33 kD. It is predominantly
	expressed on Type I IFN-producing cells (IPCs) in naive mice, but is up-regulated on most cell
	types following stimulation with type I IFNs and IFN-gamma. It is highly expressed on terminally

Target Details

differentiated normal plasmacytoid dendritic cells and some tumor cells, such as multiple myeloma, renal cell carcinoma, and melanoma cells. BST2 is a recently identified as IFN-induced cellular response factor that blocks release of HIV-1 and other retroviruses from infected cells. BST2 has been found to be the natural ligand of ILT7 in human model.

Pathways:

Regulation of Leukocyte Mediated Immunity, Production of Molecular Mediator of Immune Response


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Concentration:	0.5 mg/mL
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.