

Datasheet for ABIN2664562

anti-BST2 antibody[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	BST2
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This BST2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Clone:	927
Isotype:	IgG2b kappa
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	BST2
Alternative Name:	CD317 (BST2 Products)
Background:	CD317, known as BST2, tetherin, HM1.2 antigen, bone marrow stromal antigen 2, or PDCA-1, is type II transmembrane glycoprotein with a molecular mass of 29-33 kD. It is predominantly expressed on Type I IFN-producing cells (IPCs) in naive mice, but is up-regulated on most cell types following stimulation with type I IFNs and IFN-gamma. It is highly expressed on terminally differentiated normal plasmacytoid dendritic cells and some tumor cells, such as multiple

Target Details

myeloma, renal cell carcinoma, and melanoma cells. BST2 is a recently identified, IFN-induced cellular response factor that blocks release of HIV-1 and other retroviruses from infected cells. BST2 has been found to be the natural ligand of ILT7 in human model.

Pathways: [Regulation of Leukocyte Mediated Immunity, Production of Molecular Mediator of Immune Response](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Concentration: 0.5 mg/mL

Buffer: Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: The antibody solution should be stored undiluted between 2°C and 8°C.

Images



Flow Cytometry

Image 1.