antibodies -online.com





anti-CX3CR1 antibody





Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Overview		
Quantity:	100 μg	
Target:	CX3CR1	
Reactivity:	Mouse	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CX3CR1 antibody is un-conjugated	
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC)	
Product Details		
Clone:	SA011F11	
Isotype:	IgG2a kappa	
Purification:	The antibody was purified by affinity chromatography.	
Target Details		
Target:	CX3CR1	
Alternative Name:	CX3CR1 (CX3CR1 Products)	
Background:	CX3CR1 is a 40 kD, G-protein coupled receptor, with seven transmembrane regions. CX3CR1 is expressed by resident and alternatively activated macrophages (M2), a subset of monocytes, dendritic cells (DCs), NK cells, a subset of memory T cells, and mast cells. CX3CR1 is involved	
	in cell recruitment during inflammation and participates in cell adhesion and extravasation from	

blood vessels. Its ligand is CX3CL1, also known as fractalkine or neurotactin. CX3CR1 is also a

Target Details

coreceptor for HIV1 and variations in this gene leads to increased susceptibility to	HIV. In the
brain, it is expressed by glial cells, which interact with CX3CL1 expressed by neuro	ins.

Pathways:

Cellular Response to Molecule of Bacterial Origin


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Concentration:	0.5 mg/mL	
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.09 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	The antibody solution should be stored undiluted between 2°C and 8°C.	

Images



Flow Cytometry

Image 1.



Flow Cytometry

Image 2.