



[Go to Product page](#)

Datasheet for ABIN2666690

IL-27 Protein (AA 1-16, AA 19-228, AA 29-234)

Overview

Quantity:	10 µg
Target:	IL-27 (IL27)
Protein Characteristics:	AA 1-16, AA 19-228, AA 29-234
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Purity:	Purity is > 95 % , as determined by Coomassie stained SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 0.01 ng per µg cytokine as determined by the LAL method.

Target Details

Target:	IL-27 (IL27)
Alternative Name:	IL-27 (IL27 Products)
Background:	IL-27 is a heterodimeric cytokine that consists of EBI3, an IL-12p40-related protein, and p28, an IL-12p35-related polypeptide. IL-27 can be allocated to the IL-6/IL-12 superfamily of cytokines which include also IL-23. This family is defined based on similarities in the structural motifs of the ligands, such as a common four-helix bundle, their receptors, which contain a

Target Details

hematopoietin receptor domain. IL-27 signaling occurs via a receptor complex composed of the signal transducing receptor chains WSX-1 and glycoprotein (gp) 130. Whereas WSX-1 is the IL-27-specific receptor chain, gp130 is the common receptor subunit of IL-6 type cytokines. IL-27 signaling in naive CD4+ T cells leads to activation and phosphorylation of STAT1 and STAT3.

Molecular Weight: The total predicted molecular weight is 49 kDa. The non-reduced and the DTT-reduced proteins migrate as one band at approximately 55 kDa by SDS-PAGE.

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Comment: Biological activity: ED50 = 5 - 10 ng/ml, corresponding to a specific activity of 1.0 - 2.0 x 10⁵ units/mg, as determined by inhibition of IL-2 production induced by mouse IL-27 in mouse splenocytes activated with anti-CD3 and anti-CD28 antibodies.

Restrictions: For Research Use only

Handling

Format: Liquid

Reconstitution: For maximum results, quick spin vial prior to opening. The protein can be aliquoted and stored from -20 °C to -70 °C. Stock solutions can also be prepared at 50-100 µg/mL in sterile buffer (PBS, HPBS, DPBS, or EBSS) containing carrier protein such as 0.2-1 % BSA or HSA and stored in working aliquots at -20 °C to -70 °C.

Buffer: 0.22 µm filtered protein solution is in 20 mM NaH₂PO₄, pH 6, 0.60 M NaCl.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: -20 °C

Storage Comment: Unopened vial can be stored between 2°C and 8°C for one month, at -20°C for six months, or at -70°C for one year.