

# Datasheet for ABIN2668936 anti-STAT3 antibody (pSer727)

# 2 Images



Go to Product page

# Overview

Quantity:	200 μL
Target:	STAT3
Binding Specificity:	pSer727
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is un-conjugated
Application:	Western Blotting (WB), Dot Blot (DB)
Product Details	
Immunogen:	This STAT3 pSer727 antibody was raised against a peptide containing pSer727 of human
Immunogen:	This STAT3 pSer727 antibody was raised against a peptide containing pSer727 of human STAT3.
Immunogen: Isotype:	
	STAT3.
Isotype:	STAT3.
Isotype: Purification:	STAT3.
Isotype: Purification: Target Details	STAT3.  IgG  Affinity Purified
Isotype: Purification: Target Details Target:	STAT3.  IgG  Affinity Purified  STAT3
Isotype: Purification: Target Details Target: Alternative Name:	STAT3.  IgG  Affinity Purified  STAT3  STAT3 (STAT3 Products)

# **Target Details**

Pathways:

JAK-STAT Signaling, RTK Signaling, Interferon-gamma Pathway, Neurotrophin Signaling
Pathway, Dopaminergic Neurogenesis, Response to Growth Hormone Stimulus, Carbohydrate
Homeostasis, Stem Cell Maintenance, Hepatitis C, Protein targeting to Nucleus, Feeding
Behaviour, CXCR4-mediated Signaling Events, Signaling of Hepatocyte Growth Factor Receptor

# **Application Details**


Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze/thaw cycles and keep on ice when not in storage.
Storage:	-80 °C
Storage Comment:	Antibodies in solution can be stored at -80 °C for 2 years.
Expiry Date:	6 months

# **Images**



#### **Dot Blot**

**Image 1.** STAT3 phospho Ser727 pAb tested by dot blot analysis. Dot blot analysis was used to confirm the specificity of STAT3 phospho Ser727 pAb for STAT2 phospho Ser727. Phosphorylated peptides corresponding to the immunogen and related peptides were spotted onto PVDF and probed with the antibody at 1:30,000. The amount of peptide (picomoles) spotted is indicated next to each row. Lane 1: Unmodified Ser727 STAT1 peptide. Lane 2: Phospho Ser727 STAT1 peptide. Lane 3: Unmodified Tyr689 STAT2 peptide. Lane 4: Phospho Tyr689 STAT2



peptide. Lane 5: Unmodified Ser727 STAT3 peptide.

Lane 6: Phospho Ser727 STAT3 peptide. Lane 7:

Unmodified Tyr705 STAT3 peptide. Lane 8: Phospho

Tyr705 STAT3 peptide. Lane 9: Unmodified Ser726

STAT5A/Ser731 STAT5B peptide. Lane 10: Phospho

Ser726 STAT5A/Ser731 STAT5B peptide. Lane 11:

Unmodified Tyr694 STAT5A/Tyr699 STAT5B peptide.

Lane 12: Phospho Tyr694 STAT5A/Tyr699 STAT5B

peptide.

### **Western Blotting**

Image 2. STAT3 phospho Ser727 pAb tested by Western blot. Hep G2 nuclear extract (15 μg per lane) was probed with STAT3 phospho Ser727 pAb (1:1,000). Lane 1: cells stimulated with IL-6 (100 ng/ml for 15 minutes at 37°C). Lane 2: No treatment.