.-online.com antibodies

Datasheet for ABIN2669559 Histone H3.2 (biotinylated), (full length), (N-Term), (truncated) Protein

3 Images




Overview


Quantity:	25 μg
Target:	Histone H3.2
Protein Characteristics:	N-Term, truncated, biotinylated, full length
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	Positive Control (PC), Substrate (S)
Product Details	
Characteristics:	Truncated human Histone H3.2 is produced in E. coli and purified using FPLC. The purified protein is subsequently ligated to a peptide containing the histone tail with a N-terminal biotin via a native peptide bond. The full-length protein is then repurified prior to lyophilization. Protein concentration was determined using the molar extinction coefficient for Histone H3 and absorbance at 280nm. Protein was determined to have ≥ 98 % purity by SDS-PAGE. The molecular weight is 15,565 Daltons.
Purification:	Purified using FPLC
Purity:	Protein was determined to have \ge 98 % purity by SDS-PAGE.
Target Details	
Target:	Histone H3.2
Background:	Histone H3 is one of the core components of the nucleosome. The nucleosome is the smallest


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2669559 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	subunit of chromatin and consists of 146 base pairs of DNA wrapped around an octamer of core histone proteins (two each of H2A, H2B, H3 and H4). Histone H1 is a linker protein, present at the interface between the nucleosome core and DNA entry/exit points.
Molecular Weight:	The molecular weight is 15,565 Daltons.
Application Details	
Application Notes:	Recombinant histones are suitable for use as positive controls in the analysis of histone post- translational modifications, as substrates for histone modification enzymes, or to generate chromatin in vitro.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Recombinant histones can be resuspended in water or any suitable buffer. We recommend a starting concentration of 1 mg/mL. To fully solubilize the histone we suggest resuspension in the buffer of choice at room temperature for 20-30 minutes with occasional pipetting. Addition of salt or Tris to the resuspension buffer may enhance histone solubility.
Handling Advice:	Avoid repeated freeze/thaw cycles and keep on ice when not in storage.
Storage:	-20 °C/-80 °C
Storage Comment:	Lyophilized proteins can be stored at -20°C or -80°C, preferably desiccated. Recombinant proteins in solution are temperature sensitive and must be stored at -80°C to prevent degradation.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2669559 | 09/11/2023 | Copyright antibodies-online. All rights reserved.







Western Blotting

Image 1. Recombinant Histone H3 biotinylated tested by SDS-PAGE gel. SDS-PAGE analysis of 1.5 μ g Recombinant Histone H3 biotinylated (lane 1) and 1 μ g Recombinant Histone H3 biotinylated (lane 2). Molecular weight marker is in lane 4. Lane 3 is empty.

Mass Spectrometry

Image 2.

Mass Spectrometry

Image 3.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2669559 | 09/11/2023 | Copyright antibodies-online. All rights reserved.