antibodies

## Datasheet for ABIN2669908 Annexin V Apoptosis Detection Kit CF-Blue



Overview	
Quantity:	100 tests
Target:	Annexin V (ANXA5)
Reactivity:	Chemical
Conjugate:	CF405M
Application:	Flow Cytometry (FACS)
Product Details	
Purpose:	The kit can identify and quantitate apoptotic cells on a single-cell basis by flow cytometry. CF- Blue is an alternative to Pacific Blue dye®, BD HorizonTM V450
Sample Type:	Blood, Cell Culture Cells
Detection Method:	Fluorometric
Characteristics:	CF-Blue is an alternative to Pacific Blue dye®, BD HorizonTM V450. Staining cells simultaneously with Annexin V- CFTMBlue and the non-vital dye propidium iodide (red fluorescence) allows (bivariate analysis) the discrimination of intact cells (Annexin V- CFTMBlue negative, PI negative), early apoptotic (Annexin V- CFTMBlue positive, PI negative) and late apoptotic or necrotic cells (Annexin V- CFTMBlue positive, PI positive).
Components:	<ul> <li>100 Tests of AnnexinV CF-Blue</li> <li>100 Tests of Pl</li> <li>10X Binding Buffer</li> </ul>
Material not included:	<ul> <li>pipettes</li> <li>,</li> <li>tubes</li> </ul>

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2669908 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

## • flow cytometry machine

,

## Target Details

Target:	Annexin V (ANXA5)
Alternative Name:	Annexin V (ANXA5 Products)
Molecular Weight:	35 kDa
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Samples were tested on Flow Cytometry Induce apoptosis in cells using the desired method is not included in this time. For instance, Jurkat cells (T-cell leukemia, human) treated with 6 $\mu$ M camptothecin for four hours.
Sample Volume:	5 μL
Assay Time:	< 1 h
Protocol:	<ul> <li>Staining cells protocol with Annexin- CFTMBlue. Flow Cytometry</li> <li>1. Prepare Annexin V Binding Buffer: 10 mM Hepes/NaOH (pH 7,4) 140 mM NaCl, 2,5 mM CaCl2</li> <li>2. Induce apoptosis in cells using the desired method. A negative control should be prepared by untreated cells, that is used to define the basal level of apoptotic and necrotic or dead cells.</li> <li>3. Harvest the cells after the apoptosis induction and wash in temperate phosphate-buffered saline (PBS).</li> <li>4. Wash cells twice with temperate PBS and resuspend cells in 1 X Annexin-binding buffer at a concentration 1 x 106 cells/mL.</li> <li>5. Add 5 μL of the Annexin V- CFTMBlue and 5 μL of PI, to each 100 μL of cell suspension.</li> <li>6. Incubate the cells at room temperature for 15 minutes at room temperature (25 °C) in the dark.</li> <li>7. After incubation period, add 400 μL of 1X Annexin-binding buffer. Analyze by flow cytometry within one hour.</li> </ul>
Restrictions:	For Research Use only

## Handling

Preservative:	Sodium azide
Precaution of Use:	Reagents contain sodium azide. Sodium azide under acid conditions yields hydrazoic acid, an
	extremely toxic compound. Azide compounds should be diluted with running water before
	being discarded. These conditions are recommended to avoid deposits in plumbing where
	explosive conditions may develop.
	Do not pipet by mouth.
	Samples should be handled as if capable of transmitting infection. Appropriate disposal
	methods should be used.
	The sample preparation procedure employs a fixative (formaldehyde). Contact is to be avoided
	with skin or mucous membranes
Handling Advice:	Light exposure should be avoided. Use dim light during handling, incubation with cells and prior
	to analysis.
Storage:	4 °C