

### Datasheet for ABIN2690211

# anti-pan Keratin antibody (AA 331-430) (AbBy Fluor® 488)



	er		

Overview	
Quantity:	100 μL
Target:	pan Keratin (panKRT)
Binding Specificity:	AA 331-430
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This pan Keratin antibody is conjugated to AbBy Fluor® 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CK5

Immunogen:	KLH conjugated synthetic peptide derived from human CK5
Isotype:	IgG
Specificity:	This will target: CK4, CK75, CK3, CK2, CK6C, CK6A, keratin 6E
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse
Purification:	Purified by Protein A.

## Target Details

Target: pan Keratin (panKRT)
------------------------------

## Target Details

Alternative Name:	Keratin (panKRT Products)
Background:	Synonyms: CK4, CK75, CK3, CK2, CK6C, CK6A, keratin 6E, P-CK
	Background: Cytokeratins (CK) are intermediate filaments of epithelial cells, both in keratinising
	tissue (ie., skin) and non keratinising cells (ie., mesothelial cells). Although not a traditional
	marker for endothelial cells, cytokeratins have also been found in some microvascular
	endothelial cells. At least 20 different cytokeratins (CK) in the molecular range of 40 to 70 kDa
	and isoelectric points of 5 to 8.5 can be identified using two dimensional gel electrophoresis.
	Biochemically, most members of the CK family fall into one of two classes, type I (acidic
	polypeptides) and type II (basic polypeptides). At least one member of the acidic family and one
	member of the basic family is expressed in all epithelial cells. Defects in KRT5 are a cause of
	epidermolysis bullosa simplex.
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months