antibodies - online.com







anti-CASKIN1 antibody

Images




Overview

| 0.0 | |
|---------------------|--|
| Quantity: | 50 μg |
| Target: | CASKIN1 |
| Reactivity: | Human, Rat, Cow, Dog, Monkey, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CASKIN1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (IHC), Immunoprecipitation (IP) |
| Product Details | |
| Purification: | affinity purified |
| Target Details | |
| Target: | CASKIN1 |
| Alternative Name: | CASKIN 1 (CASKIN1 Products) |
| Application Details | |
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling


| Format: | Lyophilized |
|------------------|--|
| Reconstitution: | For reconstitution add 200 µL H2O, then aliquot and store at -20 °C until use. |
| Buffer: | PBS |
| Handling Advice: | Affinity purified antibodies are less robust than antisera, since protease inhibitors are also removed during purification. Hence, storage at 4 °C for prolonged periods (i.e. several weeks), is not recommended. |
| Storage: | -20 °C |
| Storage Comment: | Unlabeled lyophilized antibodies are stable in this form without loss of quality at ambient temperatures for several weeks or even months. They can be stored at 4°C for several years. Lyophilized antibodies must not be stored in the freezer, they may be destroyed! |

Images




Western Blotting

Image 1. dilution: 1 : 1000, sample: synaptic membrane fraction of rat brain (LP1)



Immunocytochemistry

Image 2. Indirect immunostaining of PFA fixed hippocampus neurons (dilution 1 : 500).



Immunohistochemistry

Image 3. Indirect immunostaining of a PFA fixed mouse hippocampus section (dilution 1 : 200, red).