antibodies - online.com







Mouse IL-2 ELISPOT Pair

Publications



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	5 plate
Target:	IL-2 (IL2)
Reactivity:	Mouse
Application:	ELISpot

Product Details

Brand:	BD™ ELISPOT	
Sterility:	0.2 µm filtered	
Endotoxin Level:	Endotoxin level is ≤0.01 ng/μg of protein.	
Components:	This product contains sufficient reagent for five 96-well plates, including unlabelled capture antibody (no azide/low endotoxin format), biotinylated detection antibody, and a Certificate of Analysis that provides lot-specific optimal reagent concentrations.	

Target Details

Target:	IL-2 (IL2)
Alternative Name:	IL-2 (IL2 Products)

Application Details

Application Notes:	The enzyme-linked immunospot (ELISPOT) assay is a powerful tool for detecting and
	enumerating individual cells that secrete a particular protein in vitro. Based on the sandwich
	ELISA, the ELISPOT assay derives its specificity and sensitivity by employing high affinity

capture and detection antibodies and enzyme-amplification. Although originally developed for analyzing specific antibody-secreting cells, the assay has been adapted for measuring the frequencies of cells that produce and secrete other effector molecules, such as cytokines. The sensitivity of the assay lends itself to measurement of even very low frequencies of cytokine producing cells (e.g., 1/300,000). Unique strengths of the assay include high sensitivity, high throughput, high content analysis, minimal volume of biological material required, applicability to frozen/thawed biological samples, and compatibility with other assays. For example, cells analyzed by ELISPOT can be transferred for cloning, proliferation assays, flow cytometry, or other methods of analysis.

Comment:

BD™ ELISPOT Mouse IL-2 ELISPOT Pair

Restrictions:

For Research Use only

Handling

Buffer:	No azide/low endotoxin: Aqueous buffered solution containing no preservative, 0.2µm filtered.
Preservative:	Without preservative
Storage:	4 °C
Storage Comment:	Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Publications

Product cited in:

Devin, Kassambara, Bruyer, Moreaux, Bret: "Phenotypic Characterization of Diffuse Large B-Cell Lymphoma Cells and Prognostic Impact." in: **Journal of clinical medicine**, Vol. 8, Issue 7, (2019) (PubMed).

Langer, Hammer, Hopfensperger, Klein, Hotter, De Jesus, Herbert, Pache, Smith, van der Merwe, Chanda, Fellay, Kirchhoff, Sauter: "HIV-1 Vpu is a potent transcriptional suppressor of NF-κB-elicited antiviral immune responses." in: **eLife**, Vol. 8, (2019) (PubMed).

zum Büschenfelde, Wagner, Lutzny, Oelsner, Feuerstacke, Decker, Bogner, Peschel, Ringshausen: "Recruitment of PKC-betall to lipid rafts mediates apoptosis-resistance in chronic lymphocytic leukemia expressing ZAP-70." in: **Leukemia**, Vol. 24, Issue 1, pp. 141-52, (2010) (PubMed).

Japink, Leers, Sosef, Nap: "CEA in activated macrophages. New diagnostic possibilities for

tumor markers in early colorectal cancer." in: **Anticancer research**, Vol. 29, Issue 8, pp. 3245-51, (2009) (PubMed).

El Shikh, El Sayed, Szakal, Tew: "T-independent antibody responses to T-dependent antigens: a novel follicular dendritic cell-dependent activity." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 182, Issue 6, pp. 3482-91, (2009) (PubMed).