

Datasheet for ABIN2703170

IL12 ELISA Kit





Overview

Quantity:	96 tests
Target:	IL12
Reactivity:	Rhesus Monkey
Method Type:	Sandwich ELISA
Detection Range:	40-10000 pg/mL
Minimum Detection Limit:	40 pg/mL
Application:	ELISA

Product Details

Purpose:	Rhesus Monkey IL-12 p70 ELISA Kit for Cell Culture Supernatants, Plasma, and Serum samples	
Sample Type:	Cell Culture Supernatant, Plasma, Serum	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This ELISA antibody pair detects Rhesus Monkey IL-12 p70 and human IL-12 p70. Other species not determined.	
Sensitivity:	40 pg/mL	
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data 	

Product Details

Components:

- Pre-Coated 96-well Strip Microplate
- · Wash Buffer
- · Stop Solution
- · Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

Target Details

Protein names: In 35 kDa subunit) (compared to 10 to 1		
Background: Gene Names: IL1: Protein names: In 35 kDa subunit) (0 Gene ID: 703205 UniProt: P48091 Pathways: JAK-STAT Signali Regulation of Leu		
Protein names: In 35 kDa subunit) (Gene ID: 703205 UniProt: P48091 Pathways: JAK-STAT Signali Regulation of Leu	roducts)	
UniProt: P48091 Pathways: JAK-STAT Signali Regulation of Leu	Gene Names: IL12A Protein names: Interleukin-12 subunit alpha (IL-12A) (Cytotoxic lymphocyte maturation factor 35 kDa subunit) (CLMF p35) (IL-12 subunit p35)	
Pathways: JAK-STAT Signali Regulation of Leu		
Regulation of Leu		
Activated i Cell P	ling, TLR Signaling, Cellular Response to Molecule of Bacterial Origin, ukocyte Mediated Immunity, Positive Regulation of Immune Effector Process Proliferation, Cancer Immune Checkpoints, Inflammasome	

Application Details

Application Notes:	Recommended Dilution for serum and plasma samples2 fold
Sample Volume:	100 μL
Plate:	Pre-coated

Application Details

Protocol:

- 1. Prepare all reagents, samples and standards as instructed in the manual.
- 2. Add 100 μL of standard or sample to each well.
- 3. Incubate 2.5 h at RT or O/N at 4 °C.
- 4. Add 100 μL of prepared biotin antibody to each well.
- 5. Incubate 1 h at RT.
- 6. Add 100 µL of prepared Streptavidin solution to each well.
- 7. Incubate 45 min at RT.
- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions:

For Research Use only

Handling

Storag	e.
Otorag	C.

-20 °C

Storage Comment:


The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4° C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date:

6 months

Images





Rhesus Macaque IL-12p70 concentration (pg/ml)

ELISA

Image 1.