.-online.com antibodies

Datasheet for ABIN2703440 Resistin ELISA Kit

Image



Overview

Quantity:	96 tests
Target:	Resistin (RETN)
Reactivity:	Dog
Method Type:	Sandwich ELISA
Detection Range:	16-4000 pg/mL
Minimum Detection Limit:	16 pg/mL
Application:	ELISA

Product Details

Purpose:	Canine Resistin ELISA Kit for Cell Culture Supernatants, Plasma, and Serum samples.
Sample Type:	Cell Culture Supernatant, Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Canine Resistin (RETN)
Sensitivity:	16 pg/mL
Characteristics:	 Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data
Components:	Pre-Coated 96-well Strip Microplate

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2703440 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	Wash BufferStop Solution
	Assay Diluent(s)
	Lyophilized Standard
	Biotinylated Detection Antibody
	Streptavidin-Conjugated HRP
	TMB One-Step Substrate
Material not included:	Distilled or deionized water
	- Precision pipettes to deliver 2 μ L to 1 μ L volumes
	 Adjustable 1-25 µL pipettes for reagent preparation
	 100 µL and 1 liter graduated cylinders
	Tubes to prepare standard and sample dilutions
	Absorbent paper
	Microplate reader capable of measuring absorbance at 450nm

• Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:	Resistin (RETN)
Alternative Name:	Resistin (RETN Products)
Pathways:	Feeding Behaviour, Smooth Muscle Cell Migration

Application Details

Application Notes:	Recommended Dilution for serum and plasma samples10 fold
Sample Volume:	100 µL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 μ L of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μ L of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μ L of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.


Application Details		
Restrictions:	For Research Use only	
Handling		
Storage:	-20 °C	
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is	

recommended to store at -80°C.

6 months

Expiry Date:

Images



Rat Resistin concentration (pg/ml)

