antibodies -online.com







Thyroperoxidase ELISA Kit



Image



$\overline{}$			
()	\/ \	r\/I	$\Theta \backslash \Lambda /$

Quantity:	96 tests
Target:	Thyroperoxidase (TPO)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.3-100 ng/mL
Minimum Detection Limit:	0.3 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human Thyroid Peroxidase ELISA Kit for cell culture supernatants, EDTA treated plasma, and
	serum samples. Heparin and citrate are not recommended.
Sample Type:	Plasma, Cell Culture Supernatant, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair detects human TPX. Other species not determined.
Sensitivity:	0.3 ng/mL
Characteristics:	Strip plates and additional reagents allow for use in multiple experiments
	Quantitative protein detection
	Establishes normal range
	The best products for confirmation of antibody array data

Product Details

Components:

- · Pre-Coated 96-well Strip Microplate
- Wash Buffer
- · Stop Solution
- · Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper

Thyroperoxidase (TPO)

- · Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:

Alternative Name:	Thyroid Peroxidase (TPO Products)
Background:	Gene Names: TPO
	Protein names: Thyroid peroxidase (TPO) (EC 1.11.1.8)
Gene ID:	7173
UniProt:	P07202
Pathways:	Thyroid Hormone Synthesis
Application Details	
/ Application Details	
Application Notes:	Recommended Dilution for serum and plasma samples2 fold
	Recommended Dilution for serum and plasma samples2 fold 100 µL
Application Notes:	

5. Incubate 1 h at RT.

- 6. Add 100 μL of prepared Streptavidin solution to each well.
- 7. Incubate 45 min at RT.
- 8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Sample Preparation:

Collect plasma using EDTA as an anticoagulant. Heparin and citrate are not recommended.

Restrictions:

For Research Use only

Handling

Storage:

-20 °C


Storage Comment:

The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date:

6 months

Images



ELISA

Image 1.