

Datasheet for ABIN2703508

TIMP2 ELISA Kit





Go to Product page

Overview

Quantity:	96 tests
Target:	TIMP2
Reactivity:	Human, Pig
Method Type:	Sandwich ELISA
Detection Range:	28-7.000 pg/mL
Minimum Detection Limit:	28 pg/mL
Application:	ELISA

Product Details

Purpose:	Porcine TIMP-2 ELISA Kit for cell culture supernatants, plasma, and serum samples.	
Sample Type:	Cell Culture Supernatant, Plasma, Serum	
Analytical Method:		
Detection Method:		
Specificity:	This ELISA antibody pair detects porcine and human TIMP-2. Other species not determined.	
Sensitivity:	28 pg/mL	
 Characteristics: Strip plates and additional reagents allow for use in multiple experiments Quantitative protein detection Establishes normal range The best products for confirmation of antibody array data 		
Components:	Pre-Coated 96-well Strip Microplate	

Product Details

- · Wash Buffer
- · Stop Solution
- · Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- · TMB One-Step Substrate

Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μ L to 1 μ L volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 μL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target:	TIMP2	
Alternative Name:	TIMP-2 (TIMP2 Products)	
Background:	Gene Names: TIMP-2 TIMP2 Protein names: Tissue inhibitor of metalloproteases-2	
Gene ID:	396988	
UniProt:	C0JPM4	
Pathways:	cAMP Metabolic Process	

Application Details

Application Notes:	Recommended Dilution for serum and plasma samples200 fold	
Sample Volume:	100 μL	
Plate:	Pre-coated Pre-coated	
Protocol:	 Prepare all reagents, samples and standards as instructed in the manual. Add 100 μL of standard or sample to each well. Incubate 2.5 h at RT or O/N at 4 °C. Add 100 μL of prepared biotin antibody to each well. Incubate 1 h at RT. 	

- 6. Add 100 μL of prepared Streptavidin solution to each well.
- 7. Incubate 45 min at RT.
- 8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions:

For Research Use only

Handling

Storage:	-20 °C


Storage Comment:

The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date:

6 months

Images



Porcine TIMP-2 concentration (pg/ml)

ELISA

Image 1.