antibodies -online.com







anti-CD7 antibody





Overview

Quantity:	100 μg
Target:	CD7
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD7 antibody is un-conjugated
Application:	Flow Cytometry (FACS), DNA Microscopy (DNA Mic)


Product Details

Immunogen:	T-acute lymphoblastic leukemia cells
Clone:	4H9
Isotype:	IgG2a kappa
Characteristics:	The clone 4H9 recognizes the 40 kDa CD7 antigen abundantly expressed on the human T and NK lymphocytes. The CD7 antigen is expressed throughout T lymphocyte differentiation and is present on 85 % to 90 % of peripheral blood T cells. In normal individuals, the CD7 antibody reacts with all CD8+ T cells, approximately 90 % of CD4+ T cells, and most NK cells. CD7 is weakly reactive with monocytes and does not react with granulocytes or B lymphocytes. It is also expressed on 50 % of thymocytes. In leukemias, the CD7 antigen is present on most T lymphoid lineages.
Purification:	Purified
Purity:	>95 %

Product Details Grade: **GMP** Grade **Target Details** Target: CD7 Alternative Name CD7 (CD7 Products) Background: The clone 4H9 recognizes the 40 kDa CD7 antigen abundantly expressed on the human T and NK lymphocytes. The CD7 antigen is expressed throughout T lymphocyte differentiation and is present on 85 % to 90 % of peripheral blood T cells. In normal individuals, the CD7 antibody reacts with all CD8+ T cells, approximately 90 % of CD4+ T cells, and most NK cells. CD7 is weakly reactive with monocytes and does not react with granulocytes or B lymphocytes. It is also expressed on 50 % of thymocytes. In leukemias, the CD7 antigen is present on most T lymphoid lineages. NCBI Accession: NM_006137 UniProt: P09564 Pathways: Cell-Cell Junction Organization **Application Details** Restrictions: For Research Use only Handling Format: Liquid Concentration: 1.0 mg/mL Buffer: PBS pH 7.2, 0.1 % (w/v) BSA, 0.09 % (w/v) sodium azide Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

4°C



Flow Cytometry

Image 1.

Image 2.