antibodies -online.com





DAAM1 Protein (Myc-DYKDDDDK Tag)



Image



Go to	D		
	Pron	ויאווו	mane

Overview	
Quantity:	20 μg
Target:	DAAM1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DAAM1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human DAAM1 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining


Target Details

Target:	DAAM1
Alternative Name:	Daam1 (DAAM1 Products)
Background:	Cell motility, adhesion, cytokinesis, and other functions of the cell cortex are mediated by
	reorganization of the actin cytoskeleton and several formin homology (FH) proteins have been
	associated with these processes. The protein encoded by this gene contains two FH domains
	and belongs to a novel FH protein subfamily implicated in cell polarity. A key regulator of
	cytoskeletal architecture, the small GTPase Rho, is activated during development by Wnt/Fz

Target Details

Target Details	
	signaling to control cell polarity and movement. The protein encoded by this gene is thought to function as a scaffolding protein for the Wnt-induced assembly of a disheveled (DvI)-Rho complex. This protein also promotes the nucleation and elongation of new actin filaments and regulates cell growth through the stabilization of microtubules. Alternative splicing results in multiple transcript variants encoding distinct proteins.
Molecular Weight:	123.3 kDa
NCBI Accession:	NP_055807
Pathways:	WNT Signaling
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot