# antibodies .- online.com





# Collagen Type IX Protein (Transcript Variant 2) (Myc-DYKDDDK Tag)



#### **Image**



_		_			
(40)	to I	P٢	od	uct	nad

( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Abstract:

Background:

Quantity:	20 μg	
Target:	Collagen Type IX	
Protein Characteristics:	Transcript Variant 2	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This Collagen Type IX protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	Recombinant human Collagen type IX (transcript variant 2) protein expressed in HEK293 cells.	
	Produced with end-sequenced ORF clone	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	Collagen Type IX	

This gene encodes one of the three alpha chains of type IX collagen, which is a minor (5-20 %) collagen component of hyaline cartilage. Type IX collagen is usually found in tissues containing

type II collagen, a fibrillar collagen. Studies in knockout mice have shown that synthesis of the

Collagen Type IX Products

#### **Target Details**

alpha 1 chain is essential for assembly of type IX collagen molecules, a heterotrimeric molecule, and that lack of type IX collagen is associated with early onset osteoarthritis.

Mutations in this gene are associated with osteoarthritis in humans, with multiple epiphyseal dysplasia, 6, a form of chondrodysplasia, and with Stickler syndrome, a disease characterized by ophthalmic, orofacial, articular, and auditory defects. Two transcript variants that encode different isoforms have been identified for this gene.

Molecular Weight:

62 kDa

NCBI Accession:

NP\_511040

#### **Application Details**

Application Notes: Re

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only


### Handling

Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	

immediately. Only 2-3 freeze thaw cycles are recommended.

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



## **Western Blotting**

Image 1. Validation with Western Blot